


Evidence-Based Medicine



Contents

December 2008 Vol 13 No 6

Purpose and procedure

161

EBM notebook

- 162 Jottings ...
- 162 GRADE: assessing the quality of evidence for diagnostic recommendations
- 164 Applying evidence: what's the next action?
- 166 A simple paper-based patient decision aid

Evidently

167

Therapeutics

- 168 Intensive glucose control did not prevent important complications in type 2 diabetes
- 169 Intensive glucose control increased mortality and did not prevent cardiovascular events in type 2 diabetes
- 170 Glycaemic control was not affected by self-monitoring of blood glucose in type 2 diabetes
- 171 Self-monitoring of blood glucose was not cost-effective in non-insulin-treated type 2 diabetes
- 172 Review: little evidence exists on the benefits and harms of screening for and treating gestational diabetes
- 173 Lifestyle interventions reduced the long-term risk of diabetes in adults with impaired glucose tolerance
- 174 Review: BP-lowering drugs prevent cardiovascular events equally well in younger and older patients
- 175 Metoprolol prevented myocardial infarction but increased risk of stroke and death after non-cardiac surgery
- 176 Low-carbohydrate and Mediterranean diets led to greater weight loss than a low-fat diet in moderately obese adults
- 177 Review: antioxidant supplements do not reduce all-cause mortality in primary or secondary prevention

- 178 Review: antiviral prophylaxis in pregnant women reduces recurrence of genital herpes at delivery
- 179 Review: infection rates do not differ for wounds cleansed with water or saline
- 180 Placebo acupuncture improved symptoms and quality of life in irritable bowel syndrome
- 181 An intervention promoting exclusive and prolonged breast feeding improved verbal intelligence scores in children at 6.5 years
- 182 Early, abrupt weaning did not reduce risk of HIV infection or mortality in breast-fed infants of mothers with HIV
- 183 High-concentration capsaicin reduced pain in patients with HIV-associated polyneuropathy .
- 184 Methylnaltrexone reduced opioid-induced constipation in patients with terminal illness
- 185 Early surgery was better than conservative care for short-term disability and pain in sciatica

Aetiology

- 186 Higher values of fasting plasma glucose within the normal range were associated with increased risk of type 2 diabetes

Prognosis

- 187 Review: many adults still have pain and subjective instability at 1 year after acute lateral ankle sprain

Resource reviews

188

Additional articles abstracted in ACP Journal Club


190

Other articles noted

191

Glossary

Inside back cover

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk