Evidence-Based Medicine



Contents

Purpose and procedure

EBM notebook

98 Jottings

97

- **98** Students' PEARLS: successfully incorporating evidence-based medicine in medical students' clinical attachments
- **100** The devil is in the details...or not? A primer on individual patient data meta-analysis
- **101** What kind of randomised trials do patients and clinicians need?
- **103** Selecting "Aetiology" articles for the *Evidence-Based Medicine journal*

Therapeutics

- **104** Annual screening for prostate cancer did not
- reduce mortality from prostate cancer
- **105** Periodic screening with prostatespecific antigen testing reduced mortality from prostate cancer
- **106** Review: low glycaemic-index diets reduce HbA_{1c} more than high glycaemic-index diets in diabetes mellitus
- **107** Review: multiple-session traumafocused CBT within 3 months of event reduces symptoms in acute stress disorder and PTSD
- **108** Intensive primary care treatment reduced cardiovascular risk factors in screen-detected type 2 diabetes
- **109** Review: pneumococcal vaccination is not effective for preventing pneumonia, bacteraemia, bronchitis, or mortality
- **110** Review: intranasal mupirocin ointment reduces *Staphylococcus aureus* infections in nasal carriers
- 111 Rofecoxib increased risk of CV events in patients with history of colorectal adenomas ≥1 year after stopping treatment
- **112** Oral vitamin K did not reduce bleeding in patients with excessive anticoagulation after receiving warfarin
- **113** 1 day of nitrofurantoin was not as effective as 7 days for asymptomatic bacteriuria in pregnancy
- **114** Step-up and step-down treatment strategies did not differ for symptom relief at 6 months in new onset dyspepsia

August 2009 Vol 14 No 4

- **115** Continued use of antipsychotic drugs increased longterm mortality in patients with Alzheimer disease
- **116** Review: interpersonal psychotherapy is slightly better and supportive therapy is worse than other therapies for depression
- **117** Review: back exercise interventions prevent self-reported episodes of back problems in adults but ergonomic education does not
- **118** A behavioural weight-loss programme was better than an education programme for urinary incontinence in overweight and obese women
- **119** Review: antidepressants and psychological therapies improve symptoms of irritable bowel syndrome
- **120** An exercise and telephone follow-up programme reduced emergency readmissions and improved quality of life in older people
- **121** Review: adding dexamethasone to standard therapy reduces short-term relapse for acute migraine in the emergency department

Clinical prediction guide

122 Absolute CVD risk, stratified by risk score, was 20% higher in primary care patients with CVD than in those without CVD

Quality improvement

123 Care coordination for patients with chronic conditions did not reduce hospital admissions or Medicare costs

Resource reviews

124

Additional articles abstracted in ACP Journal Club

126

Other articles noted

127

Glossary

Inside back cover

BMJIJournals

ebm.bmj.com

This article has been chosen by the Editor to

This journal is a member of and subscribes to

the principles of the Committee on Publication

www.publicationethics.org.uk

be of special interest or importance and is

freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

Ethics