Evid Based Med: first published as on 1 October 2011. Downloaded from http://ebm.bmj.com/ on April 28, 2024 by guest. Protected by copyright

Evidence-Based Medicine



Contents

Editorial

129 What is EBM?

controlled trial



Methods

Impact of facilitating physician access to relevant medical literature on outcomes of hospitalised internal medicine patients: a randomised



Prognosis

136 Thrombolysis improves 90-day functional outcome in acute ischaemic stroke, even in very older people

Therapeutics

- 138 Cardiac resynchronisation therapy is efficacious in patients with mild heart failure symptoms
- Oral rivaroxaban for acute DVT, or long term for VTE, is as effective as enoxaparin followed by a vitamin K antagonist for preventing recurrence, with no increase in bleeding complications
- 141 Lactobacillus rhamnosus GG reduces frequency and severity of abdominal pain compared with placebo in children with irritable bowel syndrome
- 142 NSAID use associated with increased cardiovascular risk and death, but naproxen appears to be the least harmful
- 143 α-blockers and antibiotics, alone or in combination, improve symptoms in men with chronic prostatitis/chronic pelvic pain syndrome
- 144 Addition of omeprazole to dual antiplatelet therapy with clopidogrel plus aspirin lowers the risk of upper gastrointestinal bleeding

October 2011 Vol 16 No 5

- 0ral antiretroviral pre-exposure prophylaxis reduces the risk of HIV acquisition among men who have sex with men
- 147 CBT stress management reduces recurrent CAD events after myocardial infarction
- 148 Initial treatment of hypertension with aliskiren and amlodipine combination gives 6.5 mm Hg greater reduction in systolic BP than does either monotherapy
- 150 Amoxicillin-clavulanate improves symptoms, reduces treatment failure in select children with acute otitis media and increases risk of diarrhoea
- **152** PREDIMED trial: Mediterranean diet may reduce the risk of type 2 diabetes
- 153 Vitamin D supplementation of deficient children may improve bone mineral density
- 154 Training birth attendants in rural Zambia in neonatal resuscitation, and the use of amoxicillin coupled with facilitated referral, reduces neonatal mortality
- Two weeks of rifaximin relieves global symptoms and bloating in people with IBS without constipation
- 157 The quadrivalent HPV vaccine is effective prophylaxis against HPV-related external genital lesions in young men
- **159** Escitalopram reduced hot flashes in non-depressed perimenopausal and postmenopausal women

Electronic page

e2 Purpose and procedure



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

