

Evidence-Based Medicine



Contents

Editorial

- 1 Evidence-based design: part of evidence-based medicine?

Perspective

- 2 Evidence-based design: medical and design researcher collaboration

Primer

- 5 Principles for high-quality, high-value testing



Systematic review

- 11 Uroradiological screening for upper and lower urinary tract anomalies in patients with hypospadias: a systematic literature review

Therapeutics

- 21 In patients with a first episode of severe alcoholic hepatitis non-responsive to medical therapy, early liver transplant increases 6-month survival
- 22 Insufficient evidence to recommend a specific bearing surface in implantable hip devices: comparative studies find little difference in outcomes
- 23 Telephone-based cognitive-behavioural therapy and a structured exercise programme are effective for chronic widespread pain (fibromyalgia)
- 25 A 10-day course of amoxicillin does not reduce symptoms of acute rhinosinusitis at day 3 of treatment
- 26 Single-tablet double-dose famotidine plus ibuprofen decreases endoscopic upper GI ulcers compared with ibuprofen alone
- 27 A small proportion of people with dementia and neuropsychiatric symptoms experience clinically significant worsening when antidepressants are discontinued
- 28 Prophylactic dose fondaparinux for 6 weeks in superficial thrombophlebitis of the legs reduces the risk for symptomatic thromboembolic complications
- 29 Oral treatment of acute pulmonary embolism with a fixed dose of rivaroxaban is non-inferior to standard treatment
- 31 Mock treatments and real acupuncture affect osteoarthritic pain similarly: patients who believe in a treatment they are receiving have less pain
- 32 Antibiotics may not improve short-term or long-term outcomes in acute uncomplicated diverticulitis
- 33 Acupuncture prophylaxis of migraine no better than sham acupuncture for decreasing frequency of headaches

February 2013 Volume 18 No 1

- 34 Closure of a patent foramen ovale with a device does not offer a greater benefit than medical therapy alone for the prevention of recurrent cerebrovascular events

Aetiology

- 36 Ranitidine use is associated with increased morbidity and mortality in very low birthweight infants
- 37 Children surviving intrauterine and neonatal insults have a significant risk of long-term adverse neurodevelopmental outcomes
- 38 The association between proton pump inhibitor use and hip fracture cannot be explained by differences in dietary and lifestyle choices

Harm

- 40 Oxycodone administered as postpartum pain relief is associated with maternal report of infant central nervous system depression in breastfed infants

Diagnosis

- 42 High-sensitivity troponin useful for diagnosis and prognosis in patients with acute coronary syndrome

Electronic pages

- e1 Brief health professional-provided interventions may lead to small improvements in physical activity
- e2 Tai chi improves balance in people with Parkinson's disease
- e3 Steroid injection and regular shoulder-specific exercises reduce the need for surgery in subacromial impingement syndrome
- e4 Dietary component of lifestyle interventions helps obese pregnant women
- e5 Family intervention focused on effective parenting is associated with decreased child obesity prevalence 3–5 years later
- e6 Primary care physicians surveyed in this study mistakenly interpreted improved survival and increased detection with screening as evidence that screening saves lives
- e7 Transition to osteoporosis for 10% of women takes 17 years for those with normal bone density, 5 years for those with moderate osteopenia: implications for the frequency of bone density testing
- e8 Emotional and physical partner abuse are common in women reporting postnatal depressive symptoms
- e9 Dabigatran associated with increased risk of acute coronary events
- e10 Patient satisfaction associated with lower emergency department use but more hospitalisations and higher costs and mortality

Receive regular table of contents by email.
Register using this QR code.



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

equator network



When you have finished with this please recycle it