


# Evidence-Based Medicine




## Contents

### Perspective

- 121** Overenthusiastic stroke risk factor modification in the over-80s: Are we being disingenuous to ourselves, and to our oldest patients?  


### Methods

- 123** Measurement of perceptions of educational environment in evidence-based medicine  


### Therapeutics

- 132** Topical antibiotic therapy is superior to systemic antibiotics for acute tympanostomy tube otorrhoea, but may not be necessary for all children
- 133** Electronic cigarettes are at least as effective as nicotine patches for smoking cessation
- 134** Cytisine is effective for smoking cessation: should clinicians use it?
- 135** Omalizumab decreases exacerbations and allows a step down in daily inhaled corticosteroid dose in adults and children with moderate-to-severe asthma
- 136** Interspinous process device versus standard conventional surgical decompression for lumbar spinal stenosis results in increased reoperation rates and costs without improving patient outcomes
- 137** Dual antiplatelet therapy in recurrent stroke prevention: do the benefits outweigh the risks?
- 138** Bariatric surgery produces greater weight loss and improvements in medical conditions than non-surgical treatment of obesity
- 139** Insufficient evidence exists to evaluate effects of blood transfusions and iron therapy for treatment of anaemia in patients with heart disease—erythropoietin agonists do not improve outcomes
- 140** The value of vitamin E as a treatment for Alzheimer's disease remains unproven despite functional improvement, due to a lack of established effect on cognition or other outcomes from RCTs
- 141** Arthroscopic partial meniscectomy provides no benefit over sham surgery in the setting of isolated degenerative medial meniscal tears without osteoarthritis
- 142** An 'all-human' diet decreases days of parenteral nutrition compared with formula in premature infants
- 143** Tight glycaemic control does not improve mortality or morbidity in critically ill children

### Prevention

- 144** No consistent evidence to date that prenatal or postnatal probiotic supplementation prevents childhood asthma and wheeze
- 145** Mailed invitations for colorectal screening are effective in increasing screening participation in uninsured adults with faecal immunochemical testing: the preferred screening modality

## August 2014 Volume 19 No 4

- 146** An intensive outpatient lifestyle modification programme improves measures of glycaemia in obese youth
- 147** Brief, patient-centred risk-reduction counselling at the time of a rapid HIV test does not affect subsequent acquisition of sexually transmitted infections
- 148** Combined HPV and cytology better than cytology for protection against cervical cancer
- 149** Extensive scheduled CT and CEA follow-up are equivalent in detecting recurrent colorectal cancer that is surgically treatable with curative intent, and superior to minimal follow up
- 150** Lung cancer screening with low-dose CT (LDCT) is ready for prime time in the USA

### Diagnosis

- 151** Voided midstream urine culture is a good test for acute cystitis in premenopausal women
- 152** Seven days of non-invasive cardiac monitoring early postischaemic stroke or TIA increases atrial fibrillation detection rate compared with current guideline-based practice

### Prognosis

- 153** Digoxin initiation is associated with lower risk of all-cause readmission after hospitalisation for heart failure
- 154** First trimester fetal size is inversely and modestly associated with cardiovascular risk factors in childhood

### Aetiology

- 155** Mediolateral episiotomy significantly reduces the risk of obstetric-associated anal sphincter injury (OASIS) in women who deliver via vacuum extraction
- 156** Migraine is a marker for risk of both ischaemic and haemorrhagic stroke
- 157** In the absence of diagnosed concussion in collegiate contact sport athletes, a relationship is suggested between the effects of head impact exposure, white matter diffusivity measures and cognition

### Quality improvement

- 158** Effect of communication training on patient, family and healthcare provider outcomes: missing links
- 159** Women with HIV infection on antiretroviral therapy with low viral loads can safely opt for vaginal delivery in the absence of obstetrical risk factors
- 160** Physician antibiotic prescriptions for skin infections in the outpatient setting are often unnecessarily long and include unnecessary antibiotics



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

[www.publicationethics.org.uk](http://www.publicationethics.org.uk)



When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

