

Evidence-Based Medicine



Contents

Perspective

- 201** Expanding the evidence within evidence-based healthcare: thinking about the context, acceptability and feasibility of interventions



Original EBM Research

- 204** Longitudinal adoption rates of complex decision support tools in primary care



Therapeutics

- 210** Intensive glucose-lowering results in increased cardiovascular mortality in younger but not older individuals with type 2 diabetes
- 211** Oseltamivir and zanamivir have limited effect on symptoms and do not reduce hospitalisation or serious complications of influenza
- 212** *Lactobacillus reuteri* is an effective option for the prevention of diarrhoea in preschool children but may not be cost-effective in all settings
- 213** Individualised pelvic floor muscle training is an effective conservative treatment in women with pelvic organ prolapse
- 214** Calcium channel blockers are effective as first line for tocolysis in the management of preterm labour
- 215** Study concludes *L. reuteri* not effective for infant colic, but findings may be limited by participants' heterogeneity
- 216** Sham treatment shows similar effects on pain and function compared to a multimodal physiotherapeutic intervention programme in patients with painful hip osteoarthritis
- 217** Induction of labour decreases a woman's chance of caesarean delivery when compared with expectant management
- 218** ACE inhibitors are associated with a reduction in all-cause mortality versus angiotensin II receptor blockers in patients with diabetes mellitus
- 219** Pro-vaccine messages may be counterproductive among vaccine-hesitant parents
- 220** At 15 years of follow-up, bariatric surgery, especially when performed within the first year, is associated with diabetes remission and reduced incidence of microvascular and macrovascular complications
- 221** Psychological therapies help reduce headache and non-headache pain in children and adolescents
- 222** Rate and rhythm control have comparable effects on mortality and stroke in atrial fibrillation but better data are needed
- 223** Ledipasvir/Sofosbuvir administration achieves very high rate of viral clearance in patients with HCV genotype 1 infection without cirrhosis, regardless of ribavirin co-administration or length of treatment

December 2014 Volume 19 No 6

- 224** Questions linger over POISE-2 and perioperative aspirin management
- 225** Surveillance for hepatocellular carcinoma with ultrasound and AFP is associated with improvements in tumour detection, receipt of curative therapy and overall survival in patients with cirrhosis
- 226** Azithromycin associated with a reduction in 90-day mortality among older pneumonia patients, although a true clinical benefit is uncertain
- 227** Until questions on intraprocedural efficacy control, renal nerve distribution and predictors of BP response are answered, the interpretation of clinical trials on renal denervation will remain uncertain
- 229** RIVUR trial offers confirmatory evidence for a small but real benefit of antibiotics for UTI prevention in children
- 230** If pharmacotherapies for alcohol use disorders are effective, why are they underutilised?

Diagnosis

- 232** Predictive values of GPs' suspicion of serious disease are high enough to warrant subsequent investigation
- 233** Improving malaria treatment by increasing access to accurate diagnostic tests: test results must guide treatment
- 234** The STONE score can help diagnose urolithiasis and decrease the use of CT scans
- 235** Prolonged cardiac monitoring after cryptogenic stroke superior to 24 h ECG in detection of occult paroxysmal atrial fibrillation

Aetiology


- 236** Elective induction of labour is associated with decreased perinatal mortality and lower odds of caesarean section at 40 and 41 weeks
- 237** Maternal body mass index increasing above 20 is associated with increased risk of miscarriage, stillbirth, neonatal death and postneonatal death


Harm

- 239** Balancing competing risks: perinatal exposure to macrolides increases the risk of infantile hypertrophic pyloric stenosis

Quality improvement

- 240** A move beyond screening is required to ensure adequate healthcare response for women who experience intimate partner violence

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk

 equator network

 recycle
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

