

Evidence-Based Medicine



Contents

August 2016 Volume 21 No 4

Editorial

- 121** Pyramids are guides not rules: the evolution of the evidence pyramid

Perspective

- 123** EBHC pyramid 5.0 for accessing preappraised evidence and guidance



EDITOR'S CHOICE

- 125** New evidence pyramid



OPEN ACCESS

- 128** Low-density lipoprotein cholesterol goals: still not in patients' best interest



- 134** Ovarian cancer screening: could you recommend it? No

Methods

- 136** Natural language processing: use in EBM and a guide for appraisal

EBM Primer

- 139** Interpreting meta-analysis of continuous outcomes with pre and postvalues

Therapeutics/Prevention

- 141** Expectant management with close monitoring of mother and neonate should be practised after late preterm prelabour rupture of membranes
- 142** Physical, nutritional, cognitive and combination interventions reduce frailty in community-dwelling prefrail and frail older people
- 143** Nebulised hypertonic saline may be less effective than previously reported in reducing hospital length of stay and admission rate in acute bronchiolitis
- 144** In *H. pylori*-infected patients second-line treatment with sequential levofloxacin therapy was more effective than levofloxacin triple therapy but was still suboptimal
- 145** Short-term health and social care benefits of the Family Nurse Partnership lack evidence in the UK context but there is promise for child developmental outcomes
- 146** Dihydroartemisinin-piperazine holds promise as an option for malaria prevention in pregnancy
- 147** Earplugs might reduce intensive care unit delirium but quality of evidence is low

- 148** Fusion in degenerative spondylolisthesis becomes controversial... again

Prognosis

- 150** Using a multiplier of 10-year cardiovascular mortality underestimates cardiovascular risk in younger individuals and women
- 151** Aortic events in patients with thoracic aortic aneurysms may be underestimated
- 152** Restarting oral anticoagulation among patients with atrial fibrillation with gastrointestinal bleeding was associated with lower risk of all-cause mortality and thromboembolism
- 153** Smoking cessation improves Crohn's disease outcomes regardless of the use of anti-TNF or immunomodulator
- 154** Risk of vascular events is lower than expected in patients with transient ischaemic attack or minor stroke treated promptly by stroke specialists
- 155** Early detection of hearing impairment reflected in better reading ability in teenage years

Aetiology/Harm

- 156** Study raises new doubts regarding the hypothesised health benefits of 'moderate' alcohol use
- 157** Study finds little impact of hydrolysed formula on risk of allergic disease, but study methods may limit conclusions and generalisability
- 158** Physicians frequently fail to de-intensify treatment in older patients with diabetes and very low haemoglobin A1c or blood pressure

Diagnosis

- 159** Large ovarian cancer screening trial shows modest mortality reduction, but does not justify population-based ovarian cancer screening

Letters

- 160** Radiation risk due to CT diagnostics of acute appendicitis
- 160** Response to letter from Professor Tapio Vehmas

This article has been chosen by the Editor to be of special interest or importance and is freely available online.

This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

C O P E
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk
Member since 2010
JM06659

equator network

recycle
When you have finished with this please recycle it

Receive regular table of contents by email. Register using this QR code.

