# **Evidence-Based Medicine**



## **Contents**

## **Editorial**

## **Primary care**

163 Redefining disease definitions and preventing overdiagnosis: time to re-evaluate our priorities

## **EBM** opinion and debate

165 Evidence-based medicine: clinicians are taught to say it but not taught to think it

## **EBM Learning**

168 Qualitative research as evidence: expanding the paradigm for evidence-based healthcare

## **EBM** analysis

#### **Primary care**



Reforming disease definitions: a new primary care led, people-centred approach

Overdiagnosis paradigm: not suitable for decreasing the overuse of antibiotics 

#### **Original EBM Research**



177 Commitments by the biopharmaceutical industry to clinical trial transparency: the evolving open access environment

## **Research methods and reporting**

## General medicine

Assessing assumptions for statistical analyses in randomised clinical trials

## **EBM Verdict**

#### **General medicine**

- Conducting one-stage IPD meta-analysis: which approach should I choose?
- Can drinking more water prevent urinary tract infections? The evidence says yes
- **192** Network meta-analysis for diagnostic tests

## October 2019 Volume 24 Issue 5

- 193 Catheter ablation in atrial fibrillation
- Routine fluoxetine in stroke

#### **Emergency care**

196 Single or dual antiplatelet therapy after a transient ischaemic attack or minor ischaemic stroke?

#### **Primary care**

- 198 Evidence from large Danish cohort does not support an association between the MMR vaccine and autism: facts in a post-truth world
- Muscle strength training for reversing frailty: how strong is the evidence?
- **200** Absolute effects of statins in the elderly

## **Electronic pages**

- Understanding Risk for Better Stroke Prevention
- e2 Pivmecillinam is more effective than ibuprofen in symptom resolution in women with uncomplicated urinary tract infection: antibiotic treatment for UTI is important
- Single-dose fosfomycin is less effective than 5-day nitrofurantoin in women with uncomplicated urinary tract infection: closing the evidence gap for a new recommendation of an old antibiotic
- Intracerebral haemorrhage is hard to stop, and must be attacked before, during and after
- CT coronary angiography does not reduce mortality or myocardial infarction in low-risk patients with acute chest pain
- **e6** Individuals with high-risk adenomas are at elevated risk for colorectal cancer
- Is there a prospect for mass drug administration of azithromycin to reduce child mortality in sub-Saharan Africa?
- e8 Biomarkers of acute kidney injury: mixed results and huge heterogeneity of reporting



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This article has been made freely available OPEN ACCESS online under the BMJ Journals open access scheme. See http://ebm.bmj.com/site/ about/guidelines.xhtml#open



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org.uk

