Evidence-Based Medicine

Contents

Purpose and procedure130	Review: tricyclic antidepressants, capsaicin, gabapentin, and oxycodone are effective for postherpetic neuralgia
Additional articles abstracted in	Review: home visiting with multidimensional assessment and multiple visits is effective in low risk elderly people 148
ACP Journal Club131 EBM notebook	Sertraline maintenance treatment reduced relapse and dropouts in post-traumatic stress disorder
Practice corner: using clinical prediction rules	Review: brief interventions reduce drinking in patients not seeking treatment
Resource review135	A brief intervention reduced alcohol drinking for up to 48 months in problem drinkers
Therapeutics	A normal calcium, low protein, low salt diet reduced recurrence of renal stones over 5 years
Oestrogen plus progestogen increased coronary heart disease and breast cancer events in postmenopausal	Physiotherapy or a wait and see policy were the best options for lateral epicondylitis at 1 year
women	Bed rest, physiotherapy, and continuation of activities of daily living were equally effective for sciatica
A lifestyle intervention or metformin prevented or delayed the onset of type 2 diabetes in people at risk	Review: antiemetic drugs reduce nausea in early pregnancy
Warfarin combined with low dose aspirin in myocardial infarction did not provide clinical benefit beyond that of	Telephone-based peer support increased the duration of breast feeding in primiparous mothers
aspirin alone	Clonidine and methylphenidate were effective for attention deficit hyperactivity disorder in children with comorbid tics
Review: angiotensin receptor blockers do not reduce mortality or hospital admission rates in heart failure	Pharmacist review changed more repeat prescriptions for
Review: n-3 polyunsaturated fatty acids reduce fatal myocardial infarction, sudden death, and overall mortality	elderly patients than usual general practice review 158
in coronary artery disease	Diagnosis
or wheeze in pre-school children was not effective 143 Review: inhaled corticosteroids for asthma or chronic	Review: questionnaires for detecting clinical depression in primary care have similar diagnostic accuracy 159
obstructive pulmonary disease do not increase bone loss 144 Review: nedocromil sodium decreases airflow reduction	Aetiology
and recovery time in exercise induced bronchoconstriction	Increased plasma total homocysteine levels increased the risk for dementia in the elderly
Review: β ₂ agonists are ineffective but increase adverse effects in acute bronchitis without underlying pulmonary disease.	Glossary Inside back cover

The virtual journal Paediatric Asthma, produced by the American Academy of Pediatrics and the BMJ Publishing Group, is now available free of charge to EBM subscribers at www.pediatric-asthma.org