


# Evidence-Based Medicine



**Editors:** Brian Haynes & Paul Glasziou

## Contents

Purpose and procedure	162		
Additional articles abstracted in <i>ACP Journal Club</i>	163		
<b>EBM notebook</b>			
Practice corner: setting EBM in motion	164		
Evidence-based decision making—the six step approach	165		
<b>Resource review</b>			
<b>Letter</b>			
Amlodipine or lisinopril was not better than chlorthalidone in lowering CHD risk in hypertension	168		
<b>Therapeutics</b>			
Review: limited evidence on regular breast examination does not support its effectiveness for reducing breast cancer deaths	169		
Oestrogen plus progestin increased risk of stroke and probable dementia in postmenopausal women	170		
Oestrogen plus progestogen increased risk of breast cancer in postmenopausal women	172		
A 10 mg nomogram was more effective than a 5 mg nomogram for warfarin induction in outpatient venous thromboembolism	173		
Lifestyle recommendations reduced blood pressure in patients with above optimal blood pressure	174		
Review: new generation antipsychotics do not induce fewer extrapyramidal side effects than low potency antipsychotics	175		
Review: ophthalmic non-steroidal anti-inflammatory drugs reduce pain without delaying healing in acute corneal abrasions	176		
Review: famotidine, pizotifen, cognitive behavioural therapy, and peppermint may be effective in recurrent abdominal pain	177		
Chemoprophylaxis with aspirin (81 mg daily) reduced the incidence of colorectal adenomas in people at risk	178		
Aspirin prevented new colorectal adenomas in patients with previous colorectal cancer	179		
		A smoking cessation intervention in a paediatric clinic setting reduced smoking in low income women	180
		An internet behavioural counselling weight loss programme reduced weight and BMI in patients at risk of type 2 diabetes	181
		Review: drugs do not improve symptoms in urinary urge incontinence and may have side effects	182
		Finasteride reduced prostate cancer but led to more high grade tumours and sexual side effects	183
		Both endurance training and strength training reduced disability and pain in chronic non-specific neck pain in women	184
		<b>Diagnosis</b>	
		The Ottawa ankle rules have a high sensitivity for excluding fractures of the ankle and midfoot in acute ankle sprain	185
		Combining fasting plasma glucose and glycosylated haemoglobin improved the accuracy for detecting patients with diabetes	186
		Conventional cervical smears were better than monolayer cytology or human papillomavirus testing for detecting cervical intraepithelial neoplasia	187
		<b>Aetiology</b>	
		Review: stimulant pharmacotherapy for children with ADHD is associated with a reduced risk of later substance abuse disorder	188
		Review: depression, social isolation, and certain life events are associated with the development of coronary heart disease	189
		<b>Quality improvement</b>	
		Review: computerised reminders and feedback can improve provider medication management	190
		Review: complex organisational and educational interventions appear to be effective for managing depression in primary care	191
		<b>Clinical prediction guide</b>	
		A simplified version of the Walsh clinical prediction rule was accurate for detecting streptococcal pharyngitis	192