

# Evidence-Based Medicine

## Contents

<b>Purpose and procedure</b> .....	162	A smoking cessation intervention in a paediatric clinic setting reduced smoking in low income women .....	180
<b>Additional articles abstracted in ACP Journal Club</b> .....	163	An internet behavioural counselling weight loss programme reduced weight and BMI in patients at risk of type 2 diabetes .....	181
<b>EBM notebook</b>		Review: drugs do not improve symptoms in urinary urge incontinence and may have side effects .....	182
Practice corner: setting EBM in motion .....	164	Finasteride reduced prostate cancer but led to more high grade tumours and sexual side effects.....	183
Evidence-based decision making—the six step approach....	165	Both endurance training and strength training reduced disability and pain in chronic non-specific neck pain in women.....	184
<b>Resource review</b> .....	167	<b>Diagnosis</b>	
<b>Letter</b>		The Ottawa ankle rules have a high sensitivity for excluding fractures of the ankle and midfoot in acute ankle sprain .....	185
Amlodipine or lisinopril was not better than chlorthalidone in lowering CHD risk in hypertension .....	168	Combining fasting plasma glucose and glycosylated haemoglobin improved the accuracy for detecting patients with diabetes .....	186
In response:.....	168	Conventional cervical smears were better than monolayer cytology or human papillomavirus testing for detecting cervical intraepithelial neoplasia.....	187
<b>Therapeutics</b>		<b>Aetiology</b>	
Review: limited evidence on regular breast examination does not support its effectiveness for reducing breast cancer deaths.....	169	Review: stimulant pharmacotherapy for children with ADHD is associated with a reduced risk of later substance abuse disorder.....	188
Oestrogen plus progestin increased risk of stroke and probable dementia in postmenopausal women.....	170	Review: depression, social isolation, and certain life events are associated with the development of coronary heart disease .....	189
Oestrogen plus progestogen increased risk of breast cancer in postmenopausal women .....	172	<b>Quality improvement</b>	
A 10 mg nomogram was more effective than a 5 mg nomogram for warfarin induction in outpatient venous thromboembolism.....	173	Review: computerised reminders and feedback can improve provider medication management .....	190
Lifestyle recommendations reduced blood pressure in patients with above optimal blood pressure .....	174	Review: complex organisational and educational interventions appear to be effective for managing depression in primary care .....	191
Review: new generation antipsychotics do not induce fewer extrapyramidal side effects than low potency antipsychotics..	175	<b>Clinical prediction guide</b>	
Review: ophthalmic non-steroidal anti-inflammatory drugs reduce pain without delaying healing in acute corneal abrasions.....	176	A simplified version of the Walsh clinical prediction rule was accurate for detecting streptococcal pharyngitis.....	192
Review: famotidine, pizotifen, cognitive behavioural therapy, and peppermint may be effective in recurrent abdominal pain .....	177	<b>Glossary</b> .....	Inside back cover
Chemoprophylaxis with aspirin (81 mg daily) reduced the incidence of colorectal adenomas in people at risk .....	178		
Aspirin prevented new colorectal adenomas in patients with previous colorectal cancer .....	179		

The virtual journal Paediatric Asthma, produced by the American Academy of Pediatrics and the BMJ Publishing Group, is now available free of charge to EBM subscribers at [www.pediatric-asthma.org](http://www.pediatric-asthma.org)